

ENTROPION AND ECTROPION CRITERIA BASED ACCESS (CBA) POLICY

| | |
|---|---|
| Version: | 2425.v1c |
| Recommendation by: | NHS Somerset ICB Clinical Commissioning Policy Forum (CCPF) |
| Date Ratified: | July 2022 |
| Name of Originator/Author: | EBI Service |
| Approved by Responsible Committee/Individual: | NHS Somerset ICB Clinical Executive Committee (CEC) |
| Publication/issue date: | September 2022 |
| Review date: | Earliest of either NICE publication or 3 years from issue |
| Target audience: | <p>NHS Somerset ICB:</p> <ul style="list-style-type: none"> • NHS Providers • GP Practices • Contracts Team <p>Medical Directors:</p> <ul style="list-style-type: none"> • Somerset Foundation Trust • Yeovil District Hospital NHS FT • Royal United Hospitals Bath NHS FT |
| Application Form | EBI Generic application form if appropriate to apply |

**ENTROPION AND ECTROPION
CRITERIA BASED ACCESS (CBA) POLICY**

| Section | CONTENTS | Page |
|----------------|--|-------------|
| | Version Control | 1 |
| 1 | General Principles | 2 |
| 2 | Policy Criteria | 3 |
| 3 | Background | 3 - 4 |
| 4 | Evidence Based Interventions Application Process | 4 - 5 |
| 5 | Access To Policy | 5 |
| 6 | References | 5 |

VERSION CONTROL

| | |
|-------------------------|----------------|
| Document Status: | Current policy |
| Version: | 2425.v1c |

DOCUMENT CHANGE HISTORY

| Version | Date | Comments |
|----------------|----------------|---|
| V1 | 2010 | Updated Guidance for Clinicians Policy Document |
| V8e | September 2015 | Reviewed by CCPF no amendments to criteria |
| 1516.v2 | July 2015 | Additional information on upper/lower lid |
| 1516.v2a | September 2015 | Trans. to CCG template, amended wording CBA lower lid / lagophthalmos |
| 1718.v4 | December 2019 | Rebranded from IFR to EBI, amended template, removal of Ectropion / Entropion & Lagophthalmos to separate policies. |
| 1920.v1 | September 2022 | 3-year review. Amendment from SCCG to NHS Somerset ICB. New PALS email address |
| 2223.v1a | February 2023 | Wording change 4.6 |
| 2223.v1b | June 2024 | Logo change with amendment to website link and clinical exceptionality wording on 4.6 |

| | |
|---|------------------|
| Equality Impact Assessment (EIA) | April 2018 |
| Quality Impact Assessment QIA | March 2018 |
| Sponsoring Director: | Dr Bernie Marden |
| Document Reference: | 2425.v1c |

1 GENERAL PRINCIPLES (CBA)

- 1.1 Treatment should only be given in line with these general principles. Where patients are unable to meet these principles, in addition to the specific treatment criteria set out in this policy, funding approval may be sought from the ICB's Evidence Based Interventions Service (EBI) by submission of an EBI application form
- 1.2 Clinicians should assess their patients against the criteria within this policy prior to a referral and/or treatment
- 1.3 Treatment should only be undertaken where the criteria have been met and there is evidence that the treatment requested is effective and the patient has the potential to benefit from the proposed treatment. Where the patient has previously been provided with the treatment with limited or diminishing benefit, it is unlikely that they will qualify for further treatment
- 1.4 Referring patients to secondary / community care without them meeting the criteria or funding approval not secured not only incurs significant costs in out-patient appointments for patients that may not qualify for surgery, but inappropriately raises the patient's expectation of treatment
- 1.5 On limited occasions, the ICB may approve funding for an assessment only in order to confirm or obtain evidence demonstrating whether a patient meets the criteria for funding. In such cases, patients should be made aware that the assessment does not mean that they will be provided with surgery and surgery will only be provided where it can be demonstrated that the patients meet the criteria to access treatment in this policy
- 1.6 Patients should be advised being referred does not confirm that they will receive treatment or surgery for a condition as a consent discussion will need to be undertaken with a clinician prior to treatment
- 1.7 The policy does not apply to patients with suspected malignancy who should continue to be referred under 2 week wait pathway rules for assessment and testing as appropriate
- 1.8 Patients with an elevated BMI of 30 or more may experience more post-surgical complications including post-surgical wound infection so should be encouraged to lose weight further prior to seeking surgery.
<https://www.sciencedirect.com/science/article/pii/S1198743X15007193>
(Thelwall, 2015)
- 1.9 Patients who are smokers should be referred to smoking cessation services in order to reduce the risk of surgery and improve healing

2 POLICY CRITERIA – CRITERIA BASED ACCESS (CBA)

2.1 Where there is a clinical concern of features suspicious of dysplasia / malignancy a referral through the local 2WW pathway should be made

2.2 ECTROPION

If a patient fulfils the criteria below no prior approval is required, please ensure the referral to secondary care indicates how the patient fulfils the criteria and the medical records evidence this

Severe Ectropion which is posing a risk to the health of the eye and conservative management has failed

2.3 ENTROPION

If a patient fulfils the criteria below no prior approval is required, please ensure the referral to secondary care indicates how the patient fulfils the criteria and the medical records evidence this

Moderate to severe Entropion where it is posing a risk to the health of the eye as eyelashes are causing persistent and on-going irritation to the eye as documented in the patient's clinical records and it is inappropriate to manage this conservatively OR

Mild Entropion, conservative management such as eye drops has failed to manage the condition and there is a significant risk to the health of the eye

2.4 Patients who are not eligible for treatment under this policy, please refer to section 4 EVIDENCE BASED INTERVENTIONS APPLICATION PROCESS on how to apply for funding with evidence of clinical exceptionality

3 BACKGROUND

3.1 Entropion

Entropion occurs when the eyelid turns inwards towards the eye. The lower lid is most commonly affected although it can rarely affect the upper lid. Entropion causes the eyelashes to rub against the front of the eye (the cornea). Severe entropion can be painful and cause vision loss by damaging the cornea. Occasionally, a corneal ulcer can form and become infected

Some eye conditions can cause this to happen, but it is also known to occur in older people associated with weakness of the small muscles around the eyelid

3.2 Ectropion

Ectropion is where the lower lid droops away from the eye and turns outwards. Most cases of ectropion are associated with ageing. They usually occur as the tissues and muscles of the eyelids become weaker as

you get older. Less common causes of ectropion include facial paralysis such as [Bell's palsy](#)

The drooping eyelid can disrupt the drainage of tears, which can make the eyes:

- Sore, red and irritated
- [Water excessively](#)
- Feel very dry and gritty
- More vulnerable to bacterial infections, such as [conjunctivitis](#)

In severe cases that aren't treated, it's possible to develop a corneal ulcer that could affect vision. However, this is rare

4 EVIDENCE BASED INTERVENTIONS APPLICATION PROCESS

4.1 Patients who are not eligible for treatment under this policy may be considered on an individual basis where their GP or Consultant believes exceptional circumstances exist that warrant deviation from the rule of this policy

4.2 Completion of a **Generic EBI Application Form** by a patient's GP or Consultant is required

4.3 Applications cannot be considered from patients personally

4.4 Only electronically completed EBI applications will be accepted to the EBI Service

4.5 It is expected that clinicians will have ensured that the patient, on behalf of who they are forwarding the application for, is appropriately informed about the existing policies prior to an application to the EBI service. This will reassure the service that the patient has a reasonable expectation of the outcome of the application and its context

4.6 EBI funding application are considered against clinical exceptionality. To eliminate discrimination for patients, **social, environmental, workplace, and non-clinical personal factors cannot be taken into consideration.**

For further information on 'clinical exceptionality' please refer to the NHS Somerset ICB EBI webpage [Evidence Based Interventions - NHS Somerset ICB](#) and click on the section titled Generic EBI Pathway.

4.7 Where appropriate photographic supporting evidence can be forwarded with the application form

4.8 An application put forward for consideration must demonstrate some unusual or unique clinical factor about the patient that suggests they are exceptional as defined below:

- Significantly different to the general population of patients with the condition in question
- Likely to gain significantly more benefit from the intervention than might be expected from the average patient with the condition

5 ACCESS TO POLICY

- 5.1 If you would like further copies of this policy or need it in another format, such as Braille or another language, please contact the Patient Advice and Liaison Service on Telephone number: 08000 851067
- 5.2 **Or write to us:** NHS Somerset ICB, Freepost RRKL-XKSC-ACSG, Yeovil, Somerset, BA22 8HR or **Email us:** somicb.pals@nhs.net

6 REFERENCES

The following sources have been considered when drafting this policy:

- 6.1 NHS Choices (2021, July)
<https://www.nhs.uk/conditions/ectropion/>
- 6.2 Orin M. Zwick, M. (2006, July). Supportive care of facial nerve palsy with temporary external eyelid weights. *Optometry - Journal of the American Optometric Association*, pp. Volume 77, Issue 7, July 2006, Pages 340–342.
- 6.3 Scott D. Lawrence, M. a. (2008, April). Lagophthalmos Evaluation and Treatment. Retrieved from America Academy of Ophthalmology:
<http://www.aao.org/eyenet/article/lagophthalmos-evaluation-treatment?april-2008>
- 6.4 Thelwall, S. P. (2015). Impact of obesity on the risk of wound infection following surgery: results from a nationwide prospective multicentre cohort study in England. *Clinical microbiology and infection : the official publication of the European Society of Clinical Microbiology and Infectious Diseases*, , vol. 21, no. 11, p. 1008.e1.