Suspected cow's milk protein allergy (CMPA) for FORMULA FED INFANTS Version 7 Clinical assessment and Family History of CMPA

Suspected mild to moderate CMPA

One or more of the following:

<u>Gastrointestinal:</u> frequent regurgitation, vomiting, diarrhoea, constipation, anaemia <u>Dermatological:</u> atopic dermatitis, urticarial (unrelated to acute infections or drugs)

Respiratory: runny nose, chronic cough, wheeze (all unrelated to infection)

General: persistent distress or colic more than 3 hours over more than 3 days over more than 3 weeks)

Formulae to trial: See prescribing guidance below.

Initial trial of Extensively hydrolysed formula (eHF) for 2-4 weeks:

- < 6 months of age: ¹Alimentum, ¹Althera, ¹Aptamil Pepti 1 or ²Nutramigen LGG 1.
- > 6 months of age: ¹Aptamil Pepti 2, ¹Alimentum, ¹Althera, or ²Nutramigen LGG 2.
- > 6 months of age: Soya infant formula Wysoy (SMA) to be purchased from supermarket/pharmacy

If weaning, will also require a strict dairy free diet. See <u>Patient webinars</u>, suitable for GPs and parents.

If no improvement after 2-4 weeks consider 2 week trial of <u>different</u> <u>eHF</u> from above list <u>before</u> considering amino acid infant formula.

THEN, If no improvement on alternative eHF, try 2-4 week trial of amino acid formula (AAF) such as Alfamino or Elecare or Nutramigen PurAmino or Neocate LCP

<u>Challenge</u>: If symptoms settle (after 4 weeks), challenge using <u>iMAP home milk challenge</u> and consider referral to Community dietitian for weaning advice.

If still symptomatic, unlikely to be only CMPA. Return to normal infant formula. Consider treating for reflux; consider referral to dietitian and/or paediatrician. **Also consider possible egg allergy.**At 1 year of age or after 6 months symptom free undertake iMAP milk challenge at home. Continue

At 1 year of age or after 6 months symptom free undertake iMAP milk challenge at home. Continue to challenge every 3 months if cow's milk still not tolerated.

At 1 year of age or after 6 months symptom free undertake milk challenge at home. See <u>Patient</u> webinars, suitable for GPs and parents. Challenge every 3 months if cow's milk still not tolerated.

Prescribing guidance

¹Alimentum, ¹Aptamil Pepti and ¹Althera- use as joint first line eHF

²Nutramigen LGG 1 & 2 and ²Neocate Syneo – <u>NOT</u> suitable for premature/immunocompromised babies

Alfamino, EleCare, Nutramigen PurAmino and Neocate LCP – Joint first line AAF

Continue infant formula as a main drink until 12 months of age

Suspected severe CMPA

One or more of the following symptoms:

<u>Gastrointestinal:</u> failure to thrive due to chronic diarrhoea, or vomiting, blood in stool, anaemia due to occult or macroscopic blood loss, protein losing enteropathy (hypoalbuminaemia), endoscopic or histologically confirmed enteropathy, severe ulcerative colitis.

<u>Dermatological:</u> urticaria, swelling (angioedema), exudative or severe atopic dermatitis

<u>Respiratory:</u> acute laryngoedema or bronchial obstruction with difficulty breathing

Systemic reactions: anaphylactic shock needs immediate hospital management EMERGENCY TREATMENT AND ADMISSION

Ongoing management: Formulae to trial: 0-12 months: Amino acid formula (AAF) e.g. Alfamino or EleCare or Nutramigen PurAmino or Neocate LCP (and strict milk free diet if weaning)

Challenge:

Home milk challenge <u>not</u> recommended.

At 12 months of age change to supermarket calcium and iodine fortified soya/oat/coconut based milk substitute.

ONLY consider prescribing eHF or AAF to a maximum of 18 months of age if CMPA **and** poor growth/intake.

May use supermarket calcium and iodine fortified soya/oat/coconut based milk substitutes in weaning foods from 6 months of age and as a main drink from 12 months of age. See patient webinars for details of appropriate products

eHF (extensively hydrolysed	formula)	AAF (Amino acid f
Alimentum (Abbott) Check availability	0-12 mths Joint 1st line; lactose free	Alfamino (Nestle)
Aptamil Pepti (Nutricia);	Contains lactose, Joint 1st line	EleCare (Abbott) Check
SMA Althera (Nestle)	Contains lactose, Joint 1st line	Nutramigen PurAming
	(may be better tolerated than Pepti due	Johnston)
	to higher hydrolysis; try before AAF	
	with continued symptoms)	
 Aptamil Pepti 1 	Joint 1st line; Under 6 months of age	Neocate (Nutricia)
 Aptamil Pepti 2 	Over 6 months of age. Joint 1st line	Neocate LCP
² Aptamil Pepti syneo	0-6 months	² Neocate Synce
Nutramigen LGG (Mead	Lactose free. Similar taste to Similac	Neocate Junior (Nutri
Johnston);	Alimentum	
 ²Nutramigen LGG 1 	Under 6 months of age	Neocate Spoon (Nutri
² Nutramigen LGG 2	Over 6 months of age	
² Nutramigen LGG 3	From 1 year (non-formulary)	
Wysoy (SMA)	May be purchased in	² Not suitable for prem
Soya based infant formula.	supermarket/pharmacy for babies over	immunocompromised
	6 months of age	

AAF (Amino acid formula)	
Alfamino (Nestle)	Joint 1 st line AAF
EleCare (Abbott) Check availability	Joint 1 st line AAF
Nutramigen PurAmino (Mead Johnston)	Joint 1 st line AAF
Neocate (Nutricia)	
 Neocate LCP 	Joint 1 st line AAF
 ²Neocate Syneo 	Non formulary
Neocate Junior (Nutricia) -	From 1 year of age. To be used under dietetic supervision only.
Neocate Spoon (Nutricia)	Specialist weaning product. To be used under dietetic supervision only.
² Not suitable for premature or immunocompromised babies	

Recomme	Amino Acid Formulae				
Products and	Alimentum £10.01	Aptamil Pepti 1	Nutramigen LGG 2	Aptamil Pepti 2	Alfamino £22.98
prices per tin	Nutramigen LGG 1 £11.21	£19.73	£11.21	£18.82	Elecare £22.98
	Aptamil Pepti 1 £9.87		Aptamil Pepti 2 £9.41		Nutramigen PurAmino £22.98
	SMA Althera £9.86				Neocate LCP £22.98
	(400g tin)	(800g tin)	(400g tin)	(800g tin)	Neocate Syneo £24.82
					(400g tin)
0 - 3 months	8	4	Not Suitable	Not Suitable	8
4 months	11	6	Not Suitable	Not Suitable	11
6 months	10	5	10	5	10
7 – 12 months	7	4	7	4	7

May use supermarket calcium and iodine fortified soya/oat/coconut based milk substitutes in weaning foods from 6 months of age and as a main drink from 12 months of age. See patient webinars for details of appropriate products.

Direct parents to patient webinars for stepping down from formula to purchased supermarket alternative at 12 months of age.

May use supermarket milk substitutes as above in <u>weaning foods</u> from 6 months of age (suitable for mild, moderate and severe CMPA) and as a main drink after 12 months of age

Milk ladder link: iMAP

https://patientwebinars.co.uk/children/children-webinars/child-allergy/

https://patientwebinars.co.uk/

NHS Somerset Infant Feeding Page