



Structured Medication Reviews on NHS Pathways

Steve Moore Locality Medicines Manager

November 28th 2023





What is an SMR?



Structured Medicine Reviews (SMRs) are an <u>evidence-based</u> and comprehensive review of a patient's medication, taking into consideration all aspects of their health. In a structured medication review clinicians and patients work as equal partners to understand the balance between the benefits and risks of and alternatives of taking medicines. The shared decision-making conversation being led by the patient's individual needs, preferences and circumstances.

- Key components of a SMR
- Shared decision-making principles should underpin the conversation
- Personalised approach tailored to the patient
- Safety consider the balance of benefit and risk of current treatment and starting new medicines
- Effectiveness all medication must be effective, noting NHS guidance on certain exceptions to items that should not routinely be prescribed in primary care."



What is NHS Pathways?



- Software we use through the Eclipse Live platform to analyse and risk stratify patients based on data extracted from the GP clinical system (emis)
- Same system that generates the red and amber patient safety alerts
- Anyone authorised can have an account to look at their own practice data
- https://secure.nhspathways.org/
- Get one time PIN
-and off you go





NHS England



Network Contract Directed Enhanced Service

Structured medication reviews and medicines optimisation: guidance

17 September 2020



Who are the focus?



From 1 October 2020, each PCN must use appropriate tools to identify and prioritise patients who would benefit from a SMR, which must include those:

- in care homes
- with complex and problematic polypharmacy, specifically those on 10 or more medications
- 3. on medicines commonly associated with medication errors 10
- with severe frailty¹¹who are particularly isolated or housebound or who have had recent hospital admissions and/or falls
- 5. using potentially addictive pain management medication.


10 See NHS Business Services Authority (2019) Medication safety – indicators specification https://www.nhsbsa.nhs.uk/sites/default/files/2019-08/Medication%20Safety%20-%20Indicators%20Specification%20%28Aug19%29.pdf

11 Based on the validation of the eFI, on average around 3% of over 65s will be identified as potentially living with severe frailty. However, this percentage may be significantly higher in some practices. Severe frailty is defined as a person having an eFI score of >0.36. https://www.england.nhs.uk/ourwork/clinical-policy/olderpeople/frailty/efi/



Landing page







Video and written guidance available



SMR Quick Search

NHS Pathways SMR Home	Patient ref:	
	Patient Code:	

Home

Data Quality Status: Extractions up to date

SMR Live Introductory Video

We are pleased to release the initial version of the SMR Live module. This is version 1.0 and we would appreciate all feedback for additional requirements.

Click here to watch our introduction video


Click here to view the SMR Live user guide

Click here to view the Structured medication reviews specification guidance 2023-24 pdf

National medicines optimisation opportunities 2023/24

This is a supplementary support tool to be utilised in conjunction with your core clinical software.







Investment and Impact Fund 22/23



A sample of pathways




•	Anticholinergic Score >= 4	717 2.4%	4 /4 8 /5	0.0% 0/717 1 / 38	- View
	Polypharmacy (>10 Meds)	850 2.9%	5 / 4 30 / 26	0.0% 0/850 1 / 38	- View
•	Pregabalin	241 0.8%	1/1 2/1	0.0% 0/241 1 / 38	- View
	Care Home	416 1.4%	1 /1 4 /4	0.0% 0/416 1 / 38	- View
•	Inhaled LABA without ICS	2 0.0%	0 /0 0 /0	0.0% 0/2 1 / 38	- View
ħ	Severe Frailty	374 1.3%	1 /1 8 /6	0.0% 0/374 1 / 38	- View
T	Learning Disability or Autism on Psychotropic drug	134 0.5%	0 / 0 0 / 0	0.0% 0/134 1 / 38	- View

National medicines optimisation opportunities 2023/24

This is a supplementary support tool to be utilised in conjunction with your core clinical software.





our

Filters: None

Core 20plus5 filter

Care home Patients: | Include



Core20PLUS5 Filter A 1erset

Age / Gender		Core20 = DD 1-2		PLUS		Vaccinations	
Age 0-17	0	Deprivation Decile 1-2	0	White	0	Flu vaccination O	
Age 18-40	0	Deprivation Decile 1-4	0	Asian	0	No flu vaccination	
Age 41-60	0	Deprivation Decile > 4	0	Black	0	All patients	
Age 61-80	0	All Deprivation Deciles	0	Ethnicity Unknown	0	Pneumococcal vaccine (last 5	
Age over 80	0			All Ethnicities	0	yrs)	
Age 65 and over	0	BMI				No Pneumococcal vaccine	
Age 18 and over	0	BMI >27.5 to 35	0	Learning Disability		(last 5 yrs)	
Age 18-60	0	BMI >35 to 40	0	Severe Mental Illness		All patients	
All Ages	0	BMI >40 to 50	0	Moderate/Severe Frailty		·	
		BMI > 50	0	Dementia		Health Check	
Male	0	All BMIs	0	Palliative Care		NHS Health Check (last 5 yrs) O	
Female	0			Depression		No NHS Health Check (last 5	
All Genders	0	Estimated Qrisk3 score		In Care Home		yrs)	
		Estimated QRISK <=10%	0	On Antipsychotics		All patients	
Smoker		Estimated QRISK3 Score >10%	. 0	On Gabapentinoids			
Current Smokers	0	Estimated QRISK3 Score >20%	, 0	•		PRISM EA	
Ex-Smoker	0	Estimated QRISK3 Score >25%	. 0	On Benzodiazepine or Z-drug		High risk O	
Current Non-Smoker	0	Estimated QRISK3 Score >30%	. 0	On Benzodiazepine		Medium risk O	
Smoking status not recorded	0	All QRISK3 Scores	0	On Z-drug		Low risk O	
All smoking statuses	0			On Opiates		No Activity	
						All patients	
*Please note that patients will need	l to fit all	conditions selected to be returned				Apply Filters	





ACB	Score: 5		
	Medication	ACB Score	
	Amitriptyline	3	✓
	Furosemide (Frusemide) Tablets	1	*
	Metoclopramide	1	*



Choose a pathway







Dependency

		Prescribing Pathway	Prevalence	Alerts	Patient Engagement	Rating
	•	Pregabalin	241 0.8%	1/1 2/1	0.0% 0/241 1 / 38	- View
	•	Gabapentin	259 0.9%	0 /0 4 /4	0.0% 0/259 1 / 38	- View
_	•	Opiates	380 1.3%	1/0 5/5	0.0% 0/380 1 / 38	- View
	•	Z-drug	188 0.6%	1/1 0/0	0.0% 0/188 1 / 38	- View
	•	Benzodiazepine	258 0.9%	0 /0 3 /2	0.0% 0/258 1 / 38	- View
	•	Smoking Protect	4,098 14.0%	1 /1 12 /10	0.0% 0/4,098 1 / 38	- View
	•	High Alcohol Intake	173 0.6%	0 /0 1 /0	0.0% 0/173 1 / 38	- View

Pathway: Z-drug ↓ NHS Somerset CCG ↓





Filters: None Core20PLUS5 Filter ➤



		Total Patients (Z-drug)	Total Patients in cohort	% Patients in cohort	
6	Moderate / Severe Frailty	187	23	12.3%	View
	Patients with morbid obesity (BMI>=40 or diabetes and BMI>=35)	187	11	5.88%	View
<u>@</u>	Current Smokers	187	39	20.86%	View
	High Alcohol intake, >= 49 units per week	187	1	0.53%	View
۵	Haemoglobin < 11	187	14	7.49%	View
•	Hyponatremia (<131)	187	3	1.6%	View
88	Polypharmacy (>10 meds)	187	42	22.46%	View
1	Patients with SMR Risk Score between 10 and 25	187	65	34.76%	View
1	Patients with SMR Risk Score > 25	187	4	2.14%	View
0	SMR: Patients with an Anticholinergic Score >= 4	187	40	21.39%	View
1	Patient in 3 SMR pathways	187	37	19.79%	View
1	Patient in 4 SMR pathways	187	18	9.63%	View
1	Patient in 5 or more SMR pathways	187	19	10.16%	View



Our Eclipse alerts



Cardiovascular	Prescribing Safety Indicators	Combined ACEI and ARB treatment MHRA warning-Combination use of medicines from different classes of reninangiotensin system blocking agents: risk of hyperkalaemia, hypotension, and impaired	26/11/2023	177	9
		renal function.https://www.gov.uk/drug-safety-update/combination-use-of-medicines- from-different-classes-of-renin-angiotensin-system-blocking-agents-risk-of- hyperkalaemia-hypotension-and-impaired-renal-function-new-warnings			
Dementia	Prescribing Safety Indicators	Dementia patients taking antipsychotic medication with no indication Increased risk of cardiovascular events and death. Should be used only after thorough review and for shortest possible time. Review regularly. Only risperidone is licensed in this condition for a maximum of 6 weeks	26/11/2023	330	1
NOACS	Prescribing Safety Indicators	DOAC patient not co-prescribed a PPI or H2RA to mitigate bleed risk Co-prescription of a PPI drug with DOACS reduces the risk of an upper GI bleed significantly. One study suggests that hospitalisation can be reduced from 4% to 2.8% by using PPIs. https://jamanetwork.com/journals/jama/fullarticle/2717474	26/11/2023	6534	32
Cardiovascular	Prescribing Safety Indicators	Glyceryl trinitrate is contraindicated in patients taking phosphodiesterase type 5 inhibitors (e.g. sildenafil, vardenafil, tadalafil) Never co-administer with: Consistent with its known effects on the nitric oxide/cyclic guanosine monophosphate (cGMP) pathway, phosphodiesterase type 5 inhibitors (e.g. sildenafil, vardenafil and tadalafil) have been shown to potentiate the hypotensive effects of nitrates (possibly resulting in collapse, unconsciousness and even death), and coadministration with glyceryl trinitrate is therefore contraindicated (see section 4.3).	26/11/2023	89	0
Other	Prescribing Safety Indicators	Learning disability on antipsychotic and no psychosis read code Learning disability on antipsychotic and no psychosis read code. NICE guidance on Psychotropic medicines in people with learning disabilities whose behaviour challenges. Use NHSE STOMP guide (stopping over medication of people with learning disabilities)to consider alternative treatments. https://www.england.nhs.uk/learning-disabilities/stomp/	26/11/2023	279	0



What practice sees



MDI/dry powder combination:	Patients 65 (Unreviewed 65)	View Patient(s)	Sun 26/11 11:23
Combined ACEI and ARB treatment:	Patients 11 (Unreviewed 11)	View Patient(s)	Sun 26/11 11:23
Carbimazole can cause congenital malformations especially at higher doses:	Patients 9 (Unreviewed 0)	View Patient(s)	Sun 26/11 11:22
Raised risk of certain skin cancers in long term hydrochlorothiazide use:	Patients 5 (Unreviewed 1)	View Patient(s)	Sun 26/11 11:20
Zonisamide in females of child bearing age:	Patients 2 (Unreviewed 0)	View Patient(s)	Sun 26/11 11:20
Severe hyperkalaemia >=6.5mmol/L:	Patients 3 (Unreviewed 3)	View Patient(s)	Sun 26/11 11:20
Warfarin in women of child bearing age not taking prescribed contraception:	Patients 7 (Unreviewed 0)	View Patient(s)	Sun 26/11 11:18
Nicorandil in active ulceration:	Patients 1 (Unreviewed 0)	View Patient(s)	Sun 26/11 11:16



If you want access...



Contact Eclipse at support@prescribingservices.org

Practice manager to send email giving consent for that person to access the practice data

Supply name, email address and mobile number (for PIN access)

Log in to https://secure.nhspathways.org/ NHS Pathways Or https://www.eclipsesolutions.org/ for Eclipse alerts

Thank you!